

UNCLASSIFIED

AD NUMBER
AD400765
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Foreign Government Information; 07 JUN 1961. Other requests shall be referred to US Library of Congress, Attn: Aerospace Technology Division, Washington, DC.
AUTHORITY
ATD ltr, 2 Dec 1965

THIS PAGE IS UNCLASSIFIED

UNCLASSIFIED

AD 400 765

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

400765

27.1.50

STEP

41386

C/002/62/028/003/001/003
F031/F004

AUTHOR: Chang, Hua (1728/5478)

TITLE: Energy band theory and the migration of excitation energy in protein molecules

PERIODICAL: Hua Hsüeh Hsüeh Pao, v. 28, no. 3, 1962, 132-138

TEXT: The theories of Szent-Gyorgyi and Schmitt and Even's calculations on the structure of proteins are either incomplete or are contradicted by experimental data. The author investigated the π -electronic structure of peptide bonds using the LCMO process and derived the energy band theory for polypeptides. Based on the protein structure theory developed by Pauling and Milliken, polypeptide molecules are considered composed of many sections and taking into account the effect of hydrogen, the author determined that monomer CO-NH has 4 π -electrons distributed on the carbonyl and amino radicals. The carbonyl radical is the electron recipient and the amino radical is the electron donor. When the radicals are put together electrons migrate from the donating radical to the receiving radical. The π -electronic systems

Card 1/9

Energy band theory and the migration ...

F031/F004

of CO-NH monomers are of three structural types: (1) Basic type — two electrons in each of the amino radicals and the carbonyl radical and the monomer is in the non-excited status. (2) Charge migration type — one π -electron of amino radical migrates to the energy level of the carbonyl radical. (3) Localized excitation type — one π -electron of carbonyl radical migrates from the radical status to the excited status. The types may be expressed in wave functions Ω_1 , Ω_2 , and Ω_3 as

$$\Omega_1 = \frac{1}{\sqrt{4!}} \sum_p (-1)^p p \{ \varphi_1(1)\alpha(1)\varphi_1(2)\beta(2)\chi_N(3)\alpha(3)\chi_N(4)\beta(4) \}$$

$$\Omega_2 = \frac{1}{\sqrt{4!}} \sum_p (-1)^p p \{ \varphi_1(1)\alpha(1)\varphi_1(2)\beta(2)\varphi_2(3)\alpha(3)\chi_N(4)\beta(4) \}$$

$$\Omega_3 = \frac{1}{\sqrt{4!}} \sum_p (-1)^p p \{ \varphi_1(1)\alpha(1)\varphi_2(2)\beta(2)\chi_N(3)\alpha(3)\chi_N(4)\beta(4) \}$$

where p is the substituting symbol; α, β are self-spinning wave functions; χ_N , the π -

Card 2/9

electron orbits of amino radical and φ_1 and φ_2 are π -electron orbits of cis- and trans- carbonyl bonds respectively. According to Kon,

$$\varphi_1 = 0.5472 X_c + 0.8370 X_o$$

$$\varphi_2 = 0.8370 X_c - 0.5472 X_o$$

where X_c and X_o are the $2p\pi$ atom orbits of carbon and oxygen respectively. The linear equation group obtained from Schrodinger's equation is

$$(H_{11} - E)C_1 + H_{12}C_2 + H_{13}C_3 = 0$$

$$H_{21}C_1 + (H_{22} - E)C_2 + H_{23}C_3 = 0$$

$$H_{31}C_1 + H_{32}C_2 + (H_{33} - E)C_3 = 0$$

where

$$H_{ii} = \int \Omega_i H \Omega_i d\tau$$

$$H_{ij} = H_{ji} = \int \Omega_j H \Omega_i d\tau$$

Card 3/9

Energy band theory and the migration ...

H is the Hamilton symbol for the CO-NH 4π -electron system. The equation group may be in the form of

$$H_{22} = I - A - (0.5472)^2 \int X_C^2(1) \left(\frac{e^2}{r_{12}}\right) X_N^2(2) d\tau_1 d\tau_2 - \\ - (0.8370)^2 \int X_O^2(1) \left(\frac{e^2}{r_{12}}\right) X_N^2(2) d\tau_1 d\tau_2$$

$$H_{12} = 0.8370 \sqrt{2} \beta_{CN}$$

$$H_{23} = -0.5472 \beta_{CN}$$

$$H_{13} = 0$$

where I is the ionization energy of the amino radical; A, the electronic attraction potential; β_{CN} , the integral of $\int X_C H X_N d\tau$. The structural wave functions Ω_1 and Ω_2 can be expressed by the following structural formulas respectively:



Card 4/9

Energy band theory and the migration ...

F031/F004

As to the status of π -electron system in peptide bond CO-NH (\dots CO-NH \dots)_{N-2} CO - NH, the above-mentioned status wave functions and corresponding energy levels satisfy equation

$$H\psi_i = E_i\psi_i, \quad i = 1, 2, 3$$

when N peptide monomers close up and form N polymers with (N-1) H bonds, the π -electron molecular orbit will change in form and the potential energy of the monomers will increase from U_n to V

$$V = \sum_{n=1}^N U_n$$

the width of energy band obeys approximately equation

$$\left(-\frac{\hbar^2}{2m} \Delta + U\right) \phi_0 = E \phi_0$$

In the N polymers, the wave function of π -electron status must satisfy Schrödinger's equation

$$\mathcal{H} \phi = \mathcal{E} \phi$$

where

$$\mathcal{H} = \left(-\frac{\hbar^2}{2m} \Delta + V\right)$$

Card 5/9

The linear combination for individual wave function of N monomers in the single peptide potential field is

$$\phi = \sum_{n=1}^N A_n \phi_0^{(n)} \quad \text{or}$$

$$\sum_{n=1}^N (\mathcal{H}_{mn} - \mathcal{J}_{mn} \mathcal{E}) A_n = 0 \quad m = 1, 2, \dots, N$$

where

$$\mathcal{H}_{mn} = \int \phi_0^{(m)} \mathcal{H} \phi_0^{(n)} d\tau$$

$$\mathcal{J}_{mn} = \int \phi_0^{(m)} \phi_0^{(n)} d\tau$$

For the same peptide monomers, \mathcal{H}_{mn} and \mathcal{J}_{mn} relate to the distance between monomers. The necessary condition for the linear equations with solutions of other than zero, is

Card 6/9

$$\begin{vmatrix}
 E - \mathcal{E} & \beta & 0 & \cdots & 0 & 0 & 0 & 0 \\
 \beta & E - \mathcal{E} & \beta & \cdots & 0 & 0 & 0 & 0 \\
 0 & \beta & E - \mathcal{E} & \cdots & 0 & 0 & 0 & 0 \\
 \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\
 0 & 0 & 0 & \cdots & E - \mathcal{E} & & \beta & \\
 0 & 0 & 0 & \cdots & \beta & & E - \mathcal{E} &
 \end{vmatrix} = 0$$

From this, we obtain energy levels

$$\mathcal{E}_m = E + 2\beta \cos \frac{m\pi}{N+1} \quad m = 1, 2, \dots, N$$

when N is large, continuous energy bands will occur, width of the band will be 4β . The difference between the the highest energy level and the lowest one is

$$W_g = E_2 - E_1 - 4|\beta|$$

Card 7/9

Take Evan's β value, the experimental value for $E_2 - E_1 = 6.1, 0.3$ from the theoretical value of 6.4. It is concluded that there are common points between cyclically and repeatedly structured polypeptides and crystals. When N monomers transit from isolated status to a regular integral body the energy levels of the monomers will split into continuous energy bands whose density is in proportion to N and whose width is in proportion to the intensity of interaction. In protein molecules the intensity is rather low, therefore the energy band is narrow (0.8 e.v.) while the width of the counter band is large (5.3 e.v.). Thus proteins are insulators under nonexcited status. However when excited, their π -electrons migrate along the hydrogen-peptide bond and form an electronic vacancy in the band and the energy of the bands is released in the form of photo-waves (fluorescence). In regularly structured protein molecules excited electrons can migrate anywhere. In the case that the order of hydrogen bonds is destroyed the common π -electron collecting energy level will not exist nor will any nonlocalized π -electron. The regularly structured protein molecules linked by hydrogen-peptide bonds may allow the migration of excited energy. The theoretical

Card 8/9

value for the migration obtained by the author is 5.3 e.v. The numerous hydrogen bonds in and among protein molecules forms the hydrogen-peptide lattice structure and the locally excited monomers enable the excited energy to migrate within the entire molecular system. Further research on the theory of the change of protein structure and the molecular vacancy caused from changes in external conditions is recommended. The author thanks Hsu Kuang-hsien (1776/0342/2009) for assistance. There is 1 figure. References in English include: A. Szent-Gyorgyi, Science 93, 609 (1941); S. Nagakura, Molecular Phys. 3, 105 (1960); K. Watanabe, J. Chem. Phys. 26, 542 (1957).

ASSOCIATION: Pei-ching Ta Hsueh Wu Li Hsi (Physics Department, Peking University)

SUBMITTED: June 7, 1961

Card 9/9